A Better Approximation Algorithm for Finding Planar Subgraphs

Gruia Călinescu * Cristina G. Fernandes † Ulrich Finkler ‡ Howard Karloff *

Abstract

The MAXIMUM PLANAR SUBGRAPH problem—given a graph G, find a largest planar subgraph of G—has applications in circuit layout, facility layout, and graph drawing. No previous polynomial-time approximation algorithm for this NP-Complete problem was known to achieve a performance ratio larger than 1/3, which is achieved simply by producing a spanning tree of G. We present the first approximation algorithm for MAXIMUM PLANAR SUBGRAPH with higher performance ratio (4/9 instead of 1/3). We also apply our algorithm to find large outerplanar subgraphs. Last, we show that both MAXIMUM PLANAR SUBGRAPH and its complement, the problem of removing as few edges as possible to leave a planar subgraph, are Max SNP-Hard.

1 Introduction

MAXIMUM PLANAR SUBGRAPH is the following problem: given a graph G, find a planar subgraph of G with the maximum number of edges. This problem has applications in circuit layout, facility layout, and graph drawing [F92, TDB88].

MAXIMUM PLANAR SUBGRAPH is known to be NP-Complete [LG77]. For a graph G, let us define $Opt(G)$ to be the maximum size of a planar subgraph of G, where *size* is the number of edges. Given an algorithm A that takes graphs G as input and outputs subgraphs of G, define $\mathrm{A}(G)$ to be the size of the planar graph A produces when G is the input. Now let us define A's *performance* or *approximation ratio* $r(\mathrm{A})$ to be the infimum, over all graphs G, of $\mathrm{A}(G)/Opt(G)$ (if $Opt(G) > 0$, and 1 otherwise). In the literature, authors sometimes ensure that their performance ratio is at least one by defining it to be the reciprocal of ours.

Numerous approximation algorithms for MAXIMUM PLANAR SUBGRAPH appear in the literature, the simplest ones being Spanning Tree (output any spanning tree of G, assuming G is connected) and Maximal Planar Subgraph (output any planar subgraph to which the addition of any new edge would violate planarity). Spanning Tree is known to have performance ratio 1/3 (see below). Dyer, Foulds and Frieze [DFF85] proved that Maximal Planar Subgraph has performance ratio 1/3. Cimikowski [Cim95] proved that a path embedding heuristic of Chiba, Nishioka and Shirakawa [CNS79] and an edge embedding heuristic of Cai, Han and Tarjan [CHT93] have performance ratios not exceeding 1/3. In the same paper, Cimikowski studied two other polynomial-time heuristics: the "vertex-addition heuristic" and the "cycle-packing heuristic." The performance ratio of the former, to the authors' knowledge, is not known, whereas for the cycle-packing algorithm, it is zero. Dyer, Foulds and Frieze [DFF85] studied two other algorithms and proved that each has performance ratio at most 2/9. Also see [JM93].

In short, to the authors' knowledge, no previously proposed algorithm was known to have a performance ratio exceeding 1/3. What makes the problem more tantalizing is that achieving a performance ratio of

*College of Computing, Georgia Institute of Technology, Atlanta, GA 30332-0280. Research supported in part by NSF grant CCR-9319106.

†College of Computing, Georgia Institute of Technology, Atlanta, GA 30332-0280. Research supported in part by the CNPq (Brazil), under contract 200975/92-7.

‡Max-Planck-Institut für Informatik, D-66123 Saarbrücken, Germany. Research supported in part by the Graduiertenkolleg Effizienz und Komplexität von Algorithmen und Rechenanlagen, Universität Saarbrücken.

1/3 is trivial. In fact, Spanning Tree has performance ratio 1/3, since every spanning tree of a connected graph on n vertices has $n-1$ edges and every planar graph on n vertices has at most $3n-3 = 3(n-1)$ edges (and there are planar graphs on n vertices with $3n-6$ edges, for all $n \geq 3$). No previous algorithm could beat the bound achieved by a trivial algorithm.

In this paper, we present two new approximation algorithms for MAXIMUM PLANAR SUBGRAPH. Each achieves a performance ratio exceeding 1/3. The higher performance ratio is $4/9 = 0.444...$ and is achieved by an algorithm which (surprisingly) invokes an algorithm for the graphic matroid parity problem as a subroutine and which runs in time $O(m^{3/2} n \log^6 n)$, where m is the number of edges in G and n is the number of vertices in G. A greedy variant still has performance ratio $7/18 = 0.3888...$, and runs in linear time on graphs of bounded degree.

Next, we provide an extension of the main algorithm. We provide a nontrivial approximation algorithm for MAXIMUM OUTERPLANAR SUBGRAPH, which is this problem: given G, find an outerplanar subgraph of G of maximum size. (An *outerplanar* graph is a graph which can be drawn in the plane without crossing edges, with all vertices on the boundary of the exterior face [H72].) This new algorithm has performance ratio at least 2/3, which surpasses the bound of 1/2 which is trivially obtained by producing a spanning tree.

Last, we show that MAXIMUM PLANAR SUBGRAPH is Max SNP-Hard, implying that there is a constant $\epsilon > 0$ such that the existence of a polynomial-time approximation algorithm with performance ratio at least $1-\epsilon$ would imply that $P = NP$ [ALMSS92]. In addition, we show that the complementary problem, called NONPLANAR DELETION or NPD—given $G = (V, E)$, produce a smallest subset $L \subseteq E$ such that $(V, E-L)$ is planar—is also Max SNP-Hard.

2 The Approximation Algorithms

In this section we present the two new algorithms for MAXIMUM PLANAR SUBGRAPH.

Let us give some motivation for our algorithm. As we said, given a (connected) graph G, an algorithm which outputs a spanning tree of G achieves a performance ratio of 1/3. A graph whose cycles all have length three, i.e., are triangles, is planar, as it cannot contain a subdivision of K_5 or $K_{3,3}$. Moreover, note that a connected spanning subgraph of G whose cycles are triangles, besides being planar, has one more edge per triangle than a spanning tree of G has.

Our better algorithm produces a subgraph of G whose cycles are triangles and, among these subgraphs, has the maximum number of edges. It can be implemented in time $O(m^{3/2} n \log^6 n)$, where m is the number of edges in G and n is the number of vertices in G, using a graphic matroid parity algorithm, as we will see later. We first present a greedy version of the algorithm.

2.1 A Greedy Version of the Algorithm

Algorithm A, presented below, is a greedy version of our algorithm. It has a performance ratio of 7/18=0.3888.... After presenting the algorithm and proving its performance ratio is 7/18, we will show it can be implemented in linear time for graphs with bounded degree. We begin with some definitions.

A *triangular cactus* is a graph whose cycles (if any) are triangles and such that all edges appear in some cycle. A *triangular cactus in a graph* G is a subgraph of G which is a triangular cactus.

A *triangular structure* is a graph whose cycles (if any) are triangles. A *triangular structure in a graph* G is a subgraph of G which is a triangular structure. Note that every triangular cactus is a triangular structure, but not *vice versa*.

Algorithm A produces a triangular structure in the given graph G. The algorithm consists of two phases. First, A greedily constructs a maximal triangular cactus S_1 in G. Second, A extends S_1 to a

triangular structure S_2 in G by adding as many edges as possible to S_1 without forming any new cycles.

Given a graph $G = (V, E)$ and $E' \subseteq E$, we denote by $G[E']$ the *spanning* subgraph of G induced by E', that is, the graph (V, E'). (Note that this is not the usual definition of subgraph induced by an edge set, since we require the subgraph to be spanning.)

Algorithm A

Starting with $E_1 = \emptyset$, repeatedly (as long as possible) find a triangle T whose vertices are in different components of $G[E_1]$, and add the edges of T to E_1.
Let $S_1 := G[E_1]$.

Starting with $E_2 = E_1$, repeatedly (as long as possible) find an edge e in G whose endpoints are in different components of $G[E_2]$, and add e to E_2.
Let $S_2 := G[E_2]$.

Output S_2.

Note that S_2 is indeed a triangular structure in G. As we mentioned before, S_2 is planar since it does not contain cycles of length greater than three.

Theorem 2.1 *The performance ratio of Algorithm A is* $\frac{7}{18}$.

Proof. First let us show that the performance ratio is at least 7/18. Without loss of generality, we may assume G is connected, and has at least three vertices. Observe that the number of edges in S_2 is the number of edges in a spanning tree of G plus the number of triangles in S_1. So it suffices to count the number of triangles in S_1.

Let H be a plane embedding of a maximum planar spanning subgraph of G. Let $n \geq 3$ be the number of vertices in G, and $t \geq 0$ be such that $3n - 6 - t$ is the number of edges in H. We can think of t as the number of edges missing for H to be a triangulated plane graph. The number of triangular faces in H is at least $2n - 4 - 2t$. (This is a lower bound on the number of triangular faces of H since if H were triangulated, it would have $2n - 4$ triangular faces, and each missing edge can destroy at most two of these triangular faces.)

Let k be the number of components of S_1 each with at least one triangle, and let $p_1, p_2, \ldots, p_k$ be the number of triangles in each of these components. Let $p = \sum_{i=1}^{k} p_i$. We will prove that p, the number of triangles in S_1, is at least a constant fraction of $n - 2 - t$. Note that if a triangle cannot be added to S_1, it is because two of its vertices are in the same component of S_1. Hence one of its edges has its two endpoints in the same component of S_1. This means that at the end of the first phase, every triangle in G must have some two vertices in the same component of S_1. In particular, every triangular face in H must have some two vertices in the same component of S_1, and therefore one of its three edges must be in the subgraph of H induced by the vertices in a component of S_1. Thus we can associate with each triangular face F in H an edge e in F whose endpoints are in the same component of S_1. But any edge e in H lies in at most two triangular faces of H, so e could have been chosen by at most two triangular faces of H. It follows that the number of triangular faces in H is at most twice the number of edges in H whose endpoints are in the same component of S_1.

Let H' be the subgraph of H induced by the edges of H whose endpoints are in the same component of S_1. Note that $p_i \geq 1$, for all i, and that the number of vertices in the i^{th} component of S_1 is $2p_i + 1 \geq 3$. Since H' is planar, H' has at most $\sum_{i=1}^{k}(3(2p_i + 1) - 6) = 6p - 3k$ edges. By the observation at the end of the previous paragraph, $2(6p - 3k) \geq 2|E(H')| \geq$ (number of triangular faces in H) $\geq 2n - 4 - 2t$. From this, we have

$$p \geq \frac{n - 2 - t + 3k}{6} \geq \frac{n - 2 - t}{6}.$$

Therefore the number of triangles in S_1 is at least $\frac{n-2-t}{6}$, and the ratio between the number of edges in S_2 and the number of edges in H is at least

$$\frac{n-1+\frac{n-2-t}{6}}{3n-6-t} = \frac{7n-8-t}{18n-36-6t} \geq \frac{7}{18},$$

since $t \geq 0$. This completes the proof that the performance ratio of Algorithm A is at least 7/18.

Now, we will prove that the performance ratio is at most 7/18. Let S be any connected triangular cactus with $p > 0$ triangles. S has $2p+1 \geq 3$ vertices. Let S' be any triangulated plane supergraph of S on the same set of vertices (S' can be obtained from S by adding edges to S until it becomes triangulated). Since S' is triangulated, S' has $2(2p+1)-4 = 4p-2$ (triangular) faces. For each face of S', add a new vertex in the face and adjacent to all vertices on the boundary of that face. Let G be the new graph. Observe that G is a triangulated plane graph and has $(2p+1)+(4p-2) = 6p-1$ vertices. This means that G has $3(6p-1)-6 = 18p-9$ edges. With G as input for Algorithm A, in the first phase it can produce $S_1 = S$, and S_2 can be S plus one edge for each of the new vertices (the vertices in G not in S). The number of edges in S is $3p$. Hence S_2 can have $3p+(4p-2) = 7p-2$ edges, while G has $18p-9$ edges. Thus the ratio between the number of edges in S_2 and the number of edges in G is

$$\frac{7p-2}{18p-9}.$$

By choosing p as large as we wish, we get a ratio as close to 7/18 as we want. ■

2.1.1 Linear Time for Bounded-Degree Graphs

In case G has bounded degree d, we can implement Algorithm A in linear time. First we describe the implementation of the first phase. At any time, the vertices of the graph are partitioned into three sets: new, active and used. At the beginning, all the vertices are new and $E_1 = \emptyset$. The following process is repeated indefinitely. Test if there are any new vertices. If there are none, halt. Test if there are any active vertices. If there are none, choose a new vertex and make it active. Choose an active vertex x and "use" it: that is, while there exist two new vertices y and z adjacent to x and to each other, choose such a pair, add the edges of the triangle xyz to E_1 and make y and z active. When there are no two new vertices adjacent to x and adjacent to each other, mark x used.

We will show in detail how "using" one vertex takes time $O(d^2)$. All the other operations—keeping lists of new and active vertices, initializing an array with the status of each vertex, initializing an auxiliary array and maintaining the list of edges in E_1—can easily be implemented in $O(n)$ time. This means the running time of this implementation is $O(nd^2)$.

Before and after "using" a vertex x, the auxiliary array has a 0 for each vertex. Going through the adjacency list of x, we change to 1 the entry of the auxiliary array for each new vertex encountered. Then, going again through the adjacency list of x, for each new vertex u encountered, go through the adjacency list of u, checking if some neighbor of u is marked 1 in the auxiliary array. This takes time $O(d)$ for each neighbor of x, for a total of $O(d^2)$ time. In this way we find, one after the other, all triangles formed by x and two new vertices. Each time such a triangle is found, mark as active the two new vertices in the triangle and place a 0 for each in the auxiliary array. At the end of this process, go through the adjacency list of x and change back to zero all remaining entries which are one in the auxiliary array.

First we note that E_1 is a triangular cactus. Indeed, at any time when we add a triangle to E_1, two of the vertices of the triangle are of type new, and since a new vertex is always an isolated vertex in $G[E_1]$, adding the three edges of the triangle keeps $G[E_1]$ a triangular cactus.

To prove that E_1 is maximal, we maintain the following invariant. All vertices which are active at a given time are in the same connected component of $G[E_1]$ at that time. To see this, note that a node u

is made active either because there are no active nodes left, in which case u is the only node active, or is made active by an active vertex x, a neighbor in G of u which is being used, and in this case the edge ux is included in $G[E_1]$.

Now, let us prove that E_1 at the end of the process is maximal, in that no triangles can be added to it. Consider a triangle xyz not in E_1. The triangle xyz has at least two vertices in the same component of $G[E_1]$ for the following reason. When the process stops, it is because there are no new vertices. Thus either x, y and z are active, or at least one of them is used. If x, y and z are active, by the invariant, the three of them are in the same component of $G[E_1]$. In the other case, assume x is the first among x, y and z to be used. While x was being used, the triangle xyz was processed. It was not added to E_1 because at least one of y and z, say y, was active. Since x was also active at that moment, x and y were (and still are) in the same component of $G[E_1]$, by the invariant.

In this paragraph we describe how to implement in linear time the second phase of Algorithm A. The time is $O(|V(G)|+|E(G)|)$ and does not depend on the graph's having bounded degree. Recall that at the end of the first phase we have a set E_1 of edges. First compute the connected components of $G[E_1]$. Let $C(v)$ be the connected component of $G[E_1]$ containing vertex $v \in V(G)$. Then construct a multigraph H in the following way: $V(H)$ is the set of connected components of $G[E_1]$, and for each edge $uv \in E(G)$, include in $E(H)$ one edge (labeled uv) with endpoints $C(u)$ and $C(v)$ if $C(u) \neq C(v)$. Run a graph transversal algorithm (for example DFS) on H and obtain a maximum spanning forest F (which we view as a subset of $E(H)$) of H. Output $E_2 = \{uv | \text{there is an edge in } F \text{ labeled } uv\}$.

2.2 A Better Algorithm

The new algorithm, Algorithm B below, finds a *maximum* triangular structure (one with the maximum number of edges) in a given graph G. Algorithm B has performance ratio 4/9, and can be implemented in time $O(m^{3/2} n \log^6 n)$. Now let us present the algorithm and the analysis of its performance ratio.

Algorithm B also has two phases. In the first one, B constructs a maximum triangular cactus S_1 in G. We will show later how to use a graphic matroid parity algorithm to construct S_1. In the second phase, B extends S_1 to a triangular structure S_2 in G, as before, by adding to S_1 as many edges as possible which do not form new cycles.

Algorithm B

Let S_1 be a *maximum* triangular cactus in G.

Starting with $E_2 = E(S_1)$, repeatedly (as long as possible) find an edge e in G whose endpoints are in different components of $G[E_2]$, and add e to E_2.

Let $S_2 := G[E_2]$.

Output S_2.

Observe that S_2 is a triangular structure in G, and therefore is planar. To analyze the algorithm, we need a definition. In any graph L, let $mts(L)$ denote the number of edges in a maximum triangular structure in L. Define $\rho(L) = mts(L)/|E(L)|$ if $E(L) \neq \emptyset$, and $\rho(L) = 1$ if $E(L) = \emptyset$.

The main result of this section is Corollary 2.11, which states that the performance ratio of Algorithm B is 4/9. As we will see, the proof of Corollary 2.11 is not difficult, given the next two theorems.

Let $G = (V, E)$ be a graph on n vertices. Let $\mathcal{P} = \{V_1, \ldots, V_k\}$ be a partition of the vertices of G and $\mathcal{Q} = \{E_1, \ldots, E_m\}$ be a partition of the edges of G. For $1 \leq i \leq m$, let u_i denote the number of classes (sometimes referred to as *color classes*) V_j of $\mathcal{P}$ met by E_i. We call the ordered pair $(\mathcal{P}, \mathcal{Q})$ *valid for* G if every triangle of G has either at least two vertices in the same part of $\mathcal{P}$ or all three edges in the same

part of $\mathcal{Q}$. Set

$$\Phi(\mathcal{P}, \mathcal{Q}) = n - k + \sum_{i=1}^{m} \lfloor \tfrac{u_i - 1}{2} \rfloor. \tag{1}$$

Note that $\Phi(\mathcal{P}, \mathcal{Q}) \geq 0$, and that there is always a valid pair $(\mathcal{P}, \mathcal{Q})$ for G (e.g., $\mathcal{P} = \{V(G)\}, \mathcal{Q} = \{E(G)\}$).

According to Lovász and Plummer [LP86], we have the following theorem:

Theorem 2.2 *The number of triangles in a maximum triangular cactus in a graph G is equal to the minimum of $\Phi(\mathcal{P}, \mathcal{Q})$ taken over all valid pairs $(\mathcal{P}, \mathcal{Q})$ for G.* [1]

Let G be a planar graph with n vertices. Embed G in the plane without crossing edges, obtaining a plane graph. Let t be the number of edges missing for this embedding to be triangulated. A triangulated plane graph has $3n - 6$ edges, if $n \geq 3$. So $t = (3n - 6) - |E(G)|$; t does not depend on the embedding.

We will prove the following theorem.

Theorem 2.3 *Let G be a connected planar graph with $n \geq 3$ vertices. Let t be the number of missing edges, defined as above. Then*

$$\Phi(\mathcal{P}, \mathcal{Q}) \geq \frac{1}{3}(n - 2 - t), \tag{2}$$

for all valid pairs $(\mathcal{P}, \mathcal{Q})$ for G.

As shown in Theorem 2.10, it follows easily from Theorems 2.2 and 2.3 that $\rho(H) \geq 4/9$ for all planar graphs H.

The proof of Theorem 2.3 is technical and long, so let us start with a general description. The proof is by induction on the number of edges in G, the basis being simple to prove, as we will see. The induction step starts with a connected plane graph G, and a valid pair $(\mathcal{P}, \mathcal{Q})$, where $\mathcal{P} = \{V_1, \ldots, V_k\}$ and $\mathcal{Q} = \{E_1, \ldots, E_m\}$. We concentrate on *singletons*, which are vertices v of G such that $\{v\} \in \mathcal{P}$.

The singletons are partitioned into three sets A, B and C as follows. Let G' be a triangulated plane supergraph of G with the same set of vertices. A is the set of singletons all of whose neighbors in G' have the same color (all neighbors in G' are in the same V_j for some j). A *facial triangle* is a triangle in G' which is the boundary of some face of G'. B is the set of singletons v not in A such that there is a facial triangle containing v with all edges in one edge class E_i for some i. We say singleton $v \in B$ *chooses* some such edge class E_i. C contains all remaining singletons.

Each singleton in B will choose an edge class E_i. For $1 \leq i \leq m$, let u_i be the number of classes V_j of $\mathcal{P}$ met by E_i, and let s_i be the number of singletons in B that have chosen E_i.

The induction step is divided into two cases. The simpler case is the one in which there is an i such that $u_i = s_i = 4$. In this case, induction is applied to a smaller graph constructed from G, and after some accounting, the theorem follows.

The hard case is the one in which there is no i such that $u_i = s_i = 4$. In this case, the result will be proved directly—induction will not be applied. Recall that t is the number of edges missing for G to be triangulated (i.e., the number of edges in G' not in G). Let s be the number of singletons, p be the number of non-singleton vertices, that is, $p = n - s$ (where n is the number of vertices of G), and q be the number of non-singleton colors, that is, $q = k - s$. Then the following upper bounds on $|A|, |B|$ and $|C|$ will be proved.

- $|A| \leq 2p - 3q - c'$ (Lemma 2.4)
- $|B| \leq 3 \sum_{i=1}^{m} \lfloor \frac{u_i - 1}{2} \rfloor$ (Lemma 2.5)

[1] Lovász was contacted and agreed that the formula given here, which differs slightly from that in [LP86], is correct.

- $|C| \leq t + c'$ (Lemma 2.6)

for some carefully chosen c'.

The upper bounds on $|A|$ and $|B|$ are not hard. The hard one is the upper bound on $|C|$. The theorem follows easily from these upper bounds and some accounting.

Proof of Theorem 2.3. By induction on the number of edges of G. Let us denote by E_G the edge set of G and by V_G its vertex set.

Basis: $|E_G| \leq 2n - 4$.

Then $t \geq (3n-6) - (2n-4) = n-2$ and therefore $\Phi(\mathcal{P}, \mathcal{Q}) \geq 0 \geq \frac{1}{3}(n-2-t)$ for all valid pairs $(\mathcal{P}, \mathcal{Q})$ for G.

Inductive step: $|E_G| > 2n - 4$.

Suppose that for any connected planar graph H on n vertices such that $|E_H| < |E_G|$, $\Phi(\mathcal{P}, \mathcal{Q}) \geq (1/3)(n - 2 - t_H)$ for all valid pairs $(\mathcal{P}, \mathcal{Q})$ for H, where t_H denotes the number of edges missing for H to be triangulated, Let us prove that $\Phi(\mathcal{P}, \mathcal{Q}) \geq (1/3)(n-2-t)$ for all valid pairs $(\mathcal{P}, \mathcal{Q})$ for G.

We begin by embedding G in the plane without crossings, obtaining a plane graph. Next, we augment G to get a triangulated plane graph G'. The edges we add are called *missing edges*. The number of missing edges in G' is t.

Let $(\mathcal{P}, \mathcal{Q})$ be a valid pair for G, with $\mathcal{P} = \{V_1, \ldots, V_k\}$ and $\mathcal{Q} = \{E_1, \ldots, E_m\}$. As before, for $1 \leq i \leq m$, let u_i denote the number of color classes V_j of $\mathcal{P}$ met by E_i. We may assume that

$$u_i \leq 2 \Rightarrow |E_i| = 1. \tag{3}$$

If $u_i \leq 2$ and $|E_i| > 1$, then we can split E_i into individual edges, obtaining a new edge partition $\mathcal{Q}'$. Any triangle in G with three edges in the old E_i also has at least two vertices of the same color, because $u_i \leq 2$. Therefore $(\mathcal{P}, \mathcal{Q}')$ is also a valid pair for G. And, moreover, $\Phi(\mathcal{P}, \mathcal{Q}') = \Phi(\mathcal{P}, \mathcal{Q})$, because $\lfloor \frac{u-1}{2} \rfloor = 0$ when $1 \leq u \leq 2$.

Assume that for $1 \leq j \leq q$, $|V_j| = p_j \geq 2$ and for $q < j \leq k$, $|V_j| = 1$. Let us call a vertex x a *singleton* if $\{x\} \in \mathcal{P}$; thus the last $k - q$ color classes are singletons. Let $s = k - q$ be the number of singletons and $p = \sum_{i=1}^{q} p_i$ be the number of vertices which are not singletons. So

$$n = p + s \quad \text{and} \quad k = q + s. \tag{4}$$

Observe that $\mathcal{P}$ is a partition of the vertices of G' (since G and G' have the same vertex set), $\mathcal{Q}$ is a partition of the edges of G (not of G'), and for any triangle of G' (not of G), facial or not, at least one of the following three conditions holds.

(1) At least two of its vertices are of the same color. We say that the triangle *is covered by a color.*

(2) All three edges are in the same class E_i of $\mathcal{Q}$. We say that the triangle *is covered by an edge class.*

(3) One or more of its edges is a missing edge. This corresponds to the fact that this triangle does not exist in G. We say that the triangle *is covered by a missing edge.*

A *facial triangle* is a triangle in G' which is the boundary of some face of G'. (Note that a facial triangle is not necessarily a triangle in G, but it is a triangle in G'.) A facial triangle T is a *facial triangle neighboring vertex* v if it contains v.

Let us partition the set of singletons into three sets A, B, C, according to how the facial triangles neighboring these singletons are covered.

A: The set of singletons all of whose neighbors in G' are of the same color.

B: The set of singletons x not in A such that there is an edge class which covers one of the facial triangles neighboring x.



Figure 1: Edges in E_i. After removing yw and, if it exists, xz, no triangles are left.

C: The set of all singletons not in $A \cup B$.

Notice that all the facial triangles neighboring a singleton in A (even the ones containing missing edges) are covered by a color. Also, not all facial triangles neighboring a singleton in C are covered by a color class (otherwise, all of its neighbors in G' would have the same color and this singleton would be in A). And no facial triangle neighboring a singleton in C is covered by an edge class (otherwise, this singleton would be put in set B, if it's not already in A).

Let a, b and c be the sizes of A, B and C, respectively. Observe that s, the number of singletons, satisfies

$$s = a + b + c. \tag{5}$$

For each singleton x in B, we choose an i such that E_i covers a facial triangle neighboring x. We denote by s_i the number of singletons x in B which have chosen E_i. Observe that E_i meets all singletons which have chosen E_i; therefore

$$s_i \leq u_i. \tag{6}$$

Moreover, if $|E_i| = 1$, then $s_i = 0$, because E_i cannot cover any triangle.

Case 1: There is an i such that $u_i = s_i = 4$.

This is the simpler case, in which we apply induction.

In this case, there are four distinct singletons x, y, w, z in B which are the only vertices E_i meets. Each of x, y, w, z has chosen E_i, implying that E_i must cover a facial triangle neighboring x. This facial triangle meets only a subset of these four vertices. Without loss of generality, we may assume this facial triangle is xyw. E_i also covers a facial triangle neighboring z. Also without loss of generality, we may assume it is ywz. Then $\{xy, yw, xw, yz, wz\} \subseteq E_i \subseteq \{xy, yw, xw, yz, wz, xz\}$ (see Figure 1).

Let G_1 be G after we remove the edges yw and xz (if edge xz exists). From the partition $\mathcal{Q}$ of $E(G)$ we obtain a partition $\mathcal{Q}_1$ of $E(G_1)$ in the following way: all parts of $\mathcal{Q}_1$ are the same as the corresponding parts of $\mathcal{Q}$, except that E_i is replaced by four parts. Each of these four parts contains exactly one edge from $E_i - \{yw, xz\}$. Note that indeed $\mathcal{Q}_1$ is a partition of $E(G_1)$. Also, the pair $(\mathcal{P}, \mathcal{Q}_1)$ is valid for G_1: each triangle in G_1 is covered exactly as it was covered in G (the triangles covered by class E_i do not exist in G_1).

Observe that G_1 is connected, and so it is a connected planar graph. Moreover, it has fewer edges than G. Thus we can apply induction, and conclude that $\Phi(\mathcal{P}, \mathcal{Q}_1) \geq \frac{1}{3}(n - 2 - t_1)$, where t_1 is the number of missing edges for G_1 to be triangulated. Because we removed at most two edges from G to get G_1, $t_1 \leq t + 2$.

The new classes in $\mathcal{Q}_1$ have zero contribution to $\Phi(\mathcal{P}, \mathcal{Q}_1)$ (since any edge class of size one contributes zero), and, since the contribution of E_i to $\Phi(\mathcal{P}, \mathcal{Q})$ is $\lfloor \frac{4-1}{2} \rfloor = 1$, it follows that $\Phi(\mathcal{P}, \mathcal{Q}_1) = \Phi(\mathcal{P}, \mathcal{Q}) - 1$.

Putting all this together, we have

$$\Phi(\mathcal{P},\mathcal{Q}) = \Phi(\mathcal{P},\mathcal{Q}_1) + 1 \geq \frac{1}{3}(n-2-t_1)+1 \geq \frac{1}{3}(n-2-(t+2))+1 > \frac{1}{3}(n-2-t).$$

Case 2: There is no i such that $u_i = s_i = 4$.

This is the hard case, whose proof requires several pages. We start by defining c', used in the upper bounds on a and c.

Consider pairs (j,F), where $1 \leq j \leq q$ and F is a face of $G'[V_j]$, the plane subgraph of G' induced by the vertex set V_j. Let c' be the number of such pairs (j,F) such that some vertex not in A is embedded in F. Call each of these c' pairs *special*.

Recall that p is the number of vertices which are not singletons, q is the number of color classes with more than one vertex, t is the number of missing edges in G', and a, b, c are the numbers of singletons in A, B, C, respectively.

The next lemmas give upper bounds on a, b and c, which will be needed to complete the proof of Theorem 2.3.

Lemma 2.4 $a \leq 2p - 3q - c'$.

Proof. Each singleton x in A has all neighbors in G' of some color $j \leq q$; hence $p_j \geq 2$. Clearly x is embedded in a face of $G'[V_G - \{x\}]$. Because all neighbors of x in G' are in V_j, this face F is also a face of $G'[V_j]$, and there cannot be any other vertex embedded in F. Therefore the size a of A is at most the number of pairs (j,F) which are not special.

The maximum number of pairs (j,F), where F is a face of $G'[V_j]$, is $2p_j - 3$ (if $p_j = 2$ then it has only one face; if $p_j > 2$ then it has the maximum number of faces when it is triangulated, and in this case, it has $2p_j - 4$ faces). Therefore there are at most $\sum_{j=1}^{q}(2p_j - 3) = 2p - 3q$ pairs (j,F), where $j \leq q$ and F is a face of $G'[V_j]$. From these pairs (j,F), c' of them are special (have a vertex which is not in A embedded in F). Thus the number of pairs (j,F) which are not special is at most $2p - 3q - c'$, and therefore $a \leq 2p - 3q - c'$. ■

Lemma 2.5 $b \leq 3\sum_{i=1}^{m} \lfloor \frac{u_i - 1}{2} \rfloor$.

Proof. If we prove that $s_i \leq 3\lfloor \frac{u_i-1}{2} \rfloor$, then by summing $s_i \leq 3\lfloor \frac{u_i-1}{2} \rfloor$ over $i \in \{1,2,...,m\}$, the lemma follows, since $b = \sum_{i=1}^{m} s_i$.

First, by (3), if $u_i \leq 2$, then $|E_i| = 1$, which implies that $s_i = 0$. And, so $s_i = 0 = 3\lfloor \frac{u_i-1}{2} \rfloor$. If $u_i > 2$, then we use the fact that inequality (6) implies that $s_i \leq u_i$. As a consequence, $s_i \leq 3\lfloor \frac{u_i-1}{2} \rfloor$ can only be false if $u_i = s_i = 4$. But this does not happen because we are given that for no i does $u_i = s_i = 4$. ■

Lemma 2.6 is the difficult upper bound. After this, we do some accounting, and the theorem follows.

Lemma 2.6 $c \leq t + c'$.

Proof. Since there is nothing to prove if $c = 0$, so we may assume $c \neq 0$, that is,

$$C \neq \emptyset. \tag{7}$$

A missing edge covers at most two facial triangles. Therefore it suffices to prove the existence of at least $2c - 2c'$ facial triangles which must be covered by missing edges.

Let us associate a set of facial triangles with each component of $G'[C]$. More specifically, for a component of $G'[C]$ whose vertex set is D, let us associate a set of either $2|D|$ or $2|D| - 2$ facial triangles, each with at

least one vertex in D, such that each such facial triangle must be covered by a missing edge. We will make sure that each of these facial triangles will have its three vertices in different color classes and at least one of them is in C. (The triangle is not covered by a color class and, because of the vertex in C, it is not covered by an edge class. So it has to be covered by a missing edge.)

Observe that the sets of facial triangles corresponding to two different components of $G'[C]$ are disjoint (since there is no edge between two different components of $G'[C]$, there cannot be a triangle with vertices in two different components of $G'[C]$). If at most c' of the components were associated with a set with $2|D|-2$ facial triangles, then in total we would have at least $2c-2c'$ facial triangles which would have to be covered by missing edges (since $\sum_D |D| = c$). This would complete the proof of Lemma 2.6.

Let D be the vertex set of a component of $G'[C]$. A *relevant facial triangle for* D is a facial triangle of G' with at least two vertices in D. Let f_D be the number of faces of $G'[D]$ (which is a connected graph), f_3 be the number of faces of $G'[D]$ which are also faces of G', and e_D be the number of edges in $G'[D]$.

Let us start by proving that $C \neq V_G$. Because $|E_G| > 2n-4$, there are at most $(3n-6)-(2n-3) = n-3$ missing edges. A missing edge covers at most two facial triangles. So there are at most $2n-6$ facial triangles covered by missing edges. But in G' there are $2n-4$ facial triangles (all the faces of G'). Thus there are facial triangles covered by colors or by edge classes. If there are facial triangles covered by colors, then at least two vertices have the same color. This means that there are vertices which are not singletons, implying that $C \neq V_G$. If all vertices are singletons and a facial triangle is covered by an edge class, then the singletons neighboring this facial triangle are in B. It follows that in either case

$$C \neq V_G. \tag{8}$$

Embedded in the faces of $G'[D]$ are all the vertices of $V_G - D$. Now $V_G - D \supseteq V_G - C \neq \emptyset$. Thus there is at least one face of $G'[D]$ which contains some vertex of $V_G - D \neq \emptyset$. This face of $G'[D]$ is not a face of G', since no face of G' contains any vertex. Thus $f_3 < f_D$.

Claim 2.7 *If $f_3 = f_D - 1$ then there are at least $2|D|-2$ relevant facial triangles for D. If $f_3 \leq f_D - 2$ then there are at least $2|D|$ relevant facial triangles for D.*

Proof. A relevant facial triangle for D has at least two vertices in D, which means it contains an edge e in $G'[D]$. But e is contained in exactly two facial triangles of G'. These are the only two relevant facial triangles for D that contain e. This would give us $2e_D$ of the desired relevant facial triangles for D, except that not all of them are distinct. If a facial triangle is counted by two edges, then all three of its vertices are in D, which implies it is a face of $G'[D]$ (and, being a facial triangle, also of G'). Moreover, it was counted exactly three times (once for each of its edges). Therefore we have $2e_D - 2f_3$ relevant facial triangles for D. Now, if $f_3 = f_D - 1$, applying Euler's formula, we get $2e_D - 2f_3 = 2e_D - 2(f_D - 1) = 2(|D|-2)+2 = 2|D|-2$ relevant facial triangles. If $f_3 \leq f_D - 2$, we get $2e_D - 2f_3 \geq 2e_D - 2(f_D - 2) = 2(|D|-2)+4 = 2|D|$. ■

Include these $2|D|$ (if possible) or $2|D|-2$ relevant facial triangles in the set of facial triangles corresponding to $G'[D]$.

To guarantee that at most c' of the components of $G'[C]$ have only $2|D|-2$ facial triangles in their corresponding sets, we will need to add two more facial triangles to some of the sets that currently have only $2|D|-2$. Now let D be the vertex set of a component of $G'[C]$ such that all the faces of $G'[D]$ but one are faces of G', that is, $f_3 = f_D - 1$. (Any component induced by a set D' with $f_3 \leq f_{D'} - 2$ already has $2|D'|$ associated facial triangles.)

Since $f_3 = f_D - 1$, all the faces of $G'[D]$ but one are faces of G'. It follows that all the vertices in $V_G - D$ must be embedded in the same face of $G'[D]$ (that face is the only face which is not a face of G'). Applying Claim 2.8 (presented below) to G' and D (as M and S, respectively), we conclude that $G'[V_G - D]$ is connected.

We use the terminology given in [BM76] for a facial walk and for the length of a face.

Claim 2.8 *Let $M = (V, E)$ be a connected plane multigraph (possibly with loops) whose faces all have length at most three. Let S be a non-empty set of vertices of M such that the subgraph $M[S]$ of M induced by S is connected, and all vertices in $V - S$ are embedded in the same face of $M[S]$. Then $M[V - S]$ is connected.*

Proof. We may assume that $V - S \neq \emptyset$. By induction on the number of edges in $M[S]$.

Basis: $M[S]$ has no edge.

$M[S]$ consists of a unique vertex u. Suppose $M[V - S]$ is not connected. Because M is connected, all the components of $M[V - S]$ are adjacent to some vertex in S, i.e., to u. The plane embedding of M determines a cyclic ordering of the edges of M incident with u. Then there exist two consecutive edges in the cyclic ordering, say e_1 and e_2, such that their other endpoints belong to different components of $M[V - S]$. But this is impossible, because then the length of the face of M lying between e_1 and e_2 would be at least four, a contradiction.

Inductive step: There is an edge in $M[S]$.

If there is an e in $M[S]$ with two distinct endpoints u and v, contract e obtaining a new connected plane multigraph H with a new vertex w corresponding to u and v. (H's vertex set V' is $(V - \{u, v\}) \cup \{w\}$.) Let $S' = (S - \{u, v\}) \cup \{w\}$. The hypotheses of the claim hold for H and S': H is connected, $H[S']$ is connected ($H[S']$ is $M[S]$ after contracting e), all the faces of H have length at most three (since contracting edges cannot increase the length of a face), $S' \neq \emptyset$, and all vertices in $V' - S' = V - S$ are embedded in the same face of $H[S']$. (There is a natural correspondence between faces of a graph and faces of the graph obtained after contracting an edge of the graph.) Furthermore, $H[S']$ has fewer edges than $M[S]$. Hence by the inductive hypotheses, $H[V' - S']$ is connected, and hence $M[V - S] = H[V' - S']$ is connected.

If every edge in $M[S]$ is a loop, then let e be such a loop, and call its unique endpoint u. Note that because $M[S]$ is connected, $S = \{u\}$. Then e divides the plane (minus e and u) into two open regions R and R'. Because all vertices of $V - S$ are embedded in the same face of $M[S]$, there is no vertex in $V - S$ embedded in one of these two regions; without loss of generality, let it be R. So there is no vertex embedded in R. This implies that any edge embedded in R is a loop, with u as its endpoint.

We consider two cases. The first case is the one in which there is some edge embedded in R. Then let H be the graph obtained from M by removing all edges of M embedded in R (they are all loops). Note that H has fewer edges than M. We will prove that the hypotheses of the claim hold for H and S. H and $H[S]$ are connected since we removed only loops. The only face of H which is not a face of M is R, and the boundary of R consists of only e: R has length one. Therefore all the faces of H have length at most three. So by the inductive hypothesis, $H[V - S]$ is connected, and hence $M[V - S] = H[V - S]$ is connected.

In the second case, there is no edge embedded in R. Hence R is a face of M (with only one edge on the boundary). Let H be the graph obtained from M by removing e. Again, H has fewer edges than M. Let F be the other face of M having e in its boundary. Removing e creates only one new face F' in H (which is the union of R, F and e). The boundary of the new face F' is the union of the boundaries of R and F minus e. The boundary of R consists of only e. Thus the boundary of F' has one edge (edge e) fewer than the boundary of F, implying that the length of F' is one less than the length of F. Therefore all the faces of H have length at most three. So by the inductive hypotheses, $H[V - S]$ is connected, and therefore $M[V - S] = H[V - S]$ is connected. ■

We proceed with the proof of Lemma 2.6.

Because $G'[D]$ is connected, all vertices of D are embedded in the same face of $G'[V_G - D]$. Call this face F'.

Claim 2.9 *One of the following holds:*

(a) *All the vertices on the boundary of F' are of the same color $j \leq q$. In this case, F' is a face of $G'[V_j]$.*
(b) *There are two distinct facial triangles of G', each with exactly one vertex in D and the other two vertices of different colors.*

Proof. The boundary of F' is a closed walk in $G'[V_G - D]$. Moreover, $V_G - D$ cannot be a set containing only one vertex x. This is true because of the following reasoning. Otherwise this vertex would have to be a singleton (because vertices in D are singletons, none can have the same color as x), more specifically, a singleton in A or in B (it could not be in C, because it is adjacent to at least one vertex of D, and hence would be in the same component of $G'[C]$ as defined by D), with all neighbors in (D and hence) C. Any vertex of G' has at least two neighbors, and a singleton x in A has all the neighbors of the same color, and therefore none of its neighbors can be a singleton. It follows that no neighbor of x can be in C; hence $x \notin A$. A singleton x in B has a neighboring facial triangle (with vertices x, y and z) covered by an edge class. But then y cannot be in C (since it has a neighboring facial triangle covered by an edge class); hence $x \notin B$.

Therefore $|V_G - D| \geq 2$. Hence the boundary of F' cannot consist of only one vertex. (If there is only one vertex on the boundary of a face of a simple plane graph, then the graph consists of only this vertex, and this does not happen because $|V_G - D| \geq 2$.)

If all vertices on the boundary of F' have the same color j, then clearly $j \leq q$ (there are at least two vertices on the boundary of F' and hence in V_j, so $j \leq q$). Also, because the only vertices embedded in F' are the vertices of D, which are not of color j, F' is also a face of $G'[V_j]$. This is case (a).

If, on the other hand, not all the vertices on the boundary of F' have the same color, then, traversing the boundary of F' (which is a closed walk), we will find at least two "oriented" edges whose endpoints have different colors. Each edge on the boundary of F' is on the boundary of exactly two triangular faces of G'. At least one of these triangular faces is contained in F'. More specifically, if an edge on the boundary of F' is traversed twice, then the two triangular faces containing this edge on the boundary are contained in F'.

Let us show that any of these triangular faces contained in F' has a vertex of D on its boundary. Suppose not. A triangular face with the three vertices on its boundary in $V_G - D$ is a face of $G'[V_G - D]$. So this triangular face would be a face of $G'[V_G - D]$ contained in F', which is a face of $G'[V_G - D]$. Because faces of a graph are disjoint, this can only happen if F' coincides with this triangular face of G'. But this is impossible, because F' is not a face of G' (all vertices in D are embedded in F').

Traversing the boundary of F', for each "oriented" edge we traverse, we find a (distinct) facial triangle of G' contained in F', containing the (unoriented version of the) oriented edge and a vertex of D on its boundary (if the two "oriented" edges are not the same unoriented edge, then each appears on the boundary of a different triangle, otherwise, the one unoriented edge appears in two distinct triangles). The facial triangles corresponding to the two "oriented" edges we picked satisfy the requirements of (b). ■

If (b) holds, add the two extra facial triangles to the set of facial triangles corresponding to $G'[D]$. Note that they are different from the ones already in the set because they contain exactly one vertex of D and all the others contain at least two vertices of D. Each set D for which (b) holds is now associated with a set of at least $2|D|$ facial triangles.

For how many components of $G'[C]$ can (a) hold? Let us show that each component of $G'[C]$ for which (a) holds corresponds to one of the c' special pairs, and that different components correspond to different special pairs.

Let D be the vertex set of a component of $G'[C]$ for which (a) holds. Let us prove that there is a special pair (j, F') such that the only vertices of C embedded in F' are the vertices in D. Let F' be the face of $G'[V_G - D]$ in which the vertices of D are embedded. Recall that $G'[V_G - D]$ is connected. Because (a) holds for this component, F' is a face of $G'[V_j]$. Since the vertices of D are embedded in F', and vertices of D are not in A, some vertex not in A is embedded in F'; this makes the pair (j, F') special. The only

vertices of G' embedded in F' are in D (because F' is a face of $G'[V_G - D]$). Since the only vertices of C embedded in F' are the vertices in D, no other component of $G'[C]$ (with a different set D) will correspond to this pair (j, F').

So the number of components such that (a) holds is at most the number c' of special pairs. Thus there are at most c' components of $G'[C]$ such that (a) holds, which completes the proof of Lemma 2.6, by the discussion in the third paragraph of the proof. ■

Using (1) and the upper bounds given by the previous lemmas, we conclude the proof of Theorem 2.3 as follows:

$$\begin{aligned}
\Phi(\mathcal{P}, \mathcal{Q}) &= n - k + \sum_{i=1}^{m} \lfloor \tfrac{u_i - 1}{2} \rfloor, \text{ by (1)}, \\
&= p - q + \sum_{i=1}^{m} \lfloor \tfrac{u_i - 1}{2} \rfloor, \text{ by (4)}, \\
&\geq p - q + \frac{1}{3} b, \text{ by Lemma 2.5}, \\
&= p - q + \frac{1}{3}(s - a - c), \text{ by Equation (5)}, \\
&\geq p - q + \frac{1}{3}(s - (2p - 3q - c') - (t + c')), \text{ by Lemmas 2.4 and 2.6}, \\
&= \frac{1}{3}(p + s) - \frac{1}{3} t \\
&= \frac{1}{3} n - \frac{1}{3} t, \text{ by the first equation in (4)}, \\
&> \frac{1}{3}(n - 2 - t).
\end{aligned}$$

This completes the proof of Theorem 2.3. ■

Recall that a triangular structure is a graph all of whose cycles are triangles. Also that $mts(L)$ denotes the number of edges in a maximum triangular structure in a graph L. And $\rho(L) = mts(L)/|E(L)|$ if $E(L) \neq \emptyset$, and $\rho(L) = 1$ if $E(L) = \emptyset$.

Theorem 2.10 *If H is a planar graph, then $\rho(H) \geq 4/9$.*

Proof. We may assume H is connected and has at least three vertices. Let t be the number of missing edges for H to be triangulated. By Euler's formula, $|E(H)| = 3n - 6 - t$, where $n \geq 3$ is the number of vertices of H.

A maximum triangular structure in H can be obtained by extending a maximum triangular cactus to a connected graph (by adding edges without forming any new cycles). Also, a maximum triangular structure in H has one more edge per triangle than a spanning tree of H.

From Theorems 2.2 and 2.3, the number of triangles in a maximum triangular cactus is at least $\frac{1}{3}(n - 2 - t)$. From this, we conclude that $mts(H) \geq n - 1 + \frac{1}{3}(n - 2 - t)$, and then

$$\rho(H) \geq \frac{n - 1 + \frac{1}{3}(n - 2 - t)}{3n - 6 - t} = \frac{4n - 5 - t}{9n - 18 - 3t} \geq \frac{4}{9},$$

for all $t \geq 0$. ■

Corollary 2.11 *The performance ratio of Algorithm B is exactly $\frac{4}{9}$.*

Proof. Let G be a graph and H be a maximum planar subgraph of G. Clearly $mts(G) \geq mts(H)$. Now $Opt(G) = |E(H)|$ implies that $\mathrm{B}(G)/Opt(G) = mts(G)/|E(H)| \geq mts(H)/|E(H)| = \rho(H)$. By Theorem 2.10, $\rho(H) \geq 4/9$ for any planar graph H. And from this we infer that the performance ratio of B is at least 4/9.

Next we prove that the performance ratio of Algorithm B is at most 4/9. Let G' be any triangulated plane graph on n' vertices. Call V' the vertex set of G'. Since G' is triangulated, G' has $2n'-4$ (triangular) faces. For each face of G', add a new vertex in the face and adjacent to all three vertices on the boundary of that face. Let G be the new graph and let V be the vertex set of G. Observe that G is a triangulated plane graph, and has $n' + (2n' - 4) = 3n' - 4$ vertices. Therefore G has $3(3n' - 4) - 6 = 9n' - 18$ edges. Let S be a maximum triangular structure in G.

Any edge in G has at least one endpoint in V'. Moreover, $|V'| = n'$. Therefore a maximum matching in G has at most n' edges (each with at least one distinct endpoint in V'). The following lemma is observed in [LP86, p. 440].

Lemma 2.12 *If S is a triangular structure with t triangles in a given graph G, then there is a matching in G of size t.*

Using Lemma 2.12, we conclude that S has at most n' triangles. Recall that S, being a triangular structure, is a spanning tree of G plus one edge per triangle in S, which implies that S has at most $(3n' - 5) + n' = 4n' - 5$ edges. Furthermore, G has $9n' - 18$ edges. Therefore the ratio between the number of edges in S and the number of edges in G is

$$\frac{4n' - 5}{9n' - 18}.$$

By choosing n' as large as we wish, we get a ratio as close to 4/9 as we want. ■

How can one find a maximum triangular cactus quickly? A graphic matroid parity algorithm can be used to construct a maximum triangular cactus in a given graph [LP86]. The problem solved by a graphic matroid parity algorithm is GRAPHIC MATROID PARITY (GMP): given a multigraph $G' = (V', E')$ and a partition of the edge set E' into pairs of distinct edges $\{f, f'\}$, find a (simple) forest F with the maximum number of edges, such that $f \in F$ if and only if $f' \in F$, for all $f \in E'$.

Let us show how to reduce the problem of finding a maximum triangular cactus in a given graph $G = (V, E)$ to GMP. This is done by describing a multigraph $G' = (V', E')$ and a partition $\mathcal{P}$ of E' into pairs of distinct edges of E', such that, from a solution to GMP for G' and $\mathcal{P}$, we can construct a maximum triangular cactus in G.

First, let $V' = V$. Now, let us describe E' and the partition $\mathcal{P}$. Initially, $E' = \emptyset$ and $\mathcal{P} = \emptyset$. For each triangle in G with edge set T, let e, e' be any pair of distinct edges in T. Add two new edges f and f' to E', f with the same endpoints as e, and f' with the same endpoints as e'. We say that T *corresponds to* $\{f, f'\}$. Insert $\{f, f'\}$ into $\mathcal{P}$. The construction of E' and $\mathcal{P}$ is complete.

We say a forest F in G' is *valid* if $f \in F$ if and only if $f' \in F$, for all f in E'. Observe that any valid forest has an even number of edges. The following lemma states a relation between valid forests in G' and triangular cacti in G. Let m and n be the number of edges and vertices, respectively, in G.

Lemma 2.13 *There is a valid forest F in G' with $2p$ edges if and only if there is a triangular cactus S in G with p triangles. Moreover, S can be obtained from F in time $O(n)$.*

Proof. Assume that S is a triangular cactus in G with p triangles. Note that an edge in S appears in exactly one cycle; otherwise that edge is a chord in a cycle of length at least four. So if in each triangle we replace the three edges by the two edges f, f', we are left with a valid forest in G', with $2p$ edges.

Assume now that F is a valid forest in G' with $2p$ edges. Note that $2p \leq n-1$. For each pair $\{f, f'\}$ in F, substitute f, f' by e, e' and add the third edge of the corresponding triangle. The substitution can be done in $O(1)$ time per pair, in a total time of $O(p)$, which is $O(n)$.

Call the resulting graph S. Next we prove that S is a triangular cactus with p triangles. Note that S is a subgraph of G. Each edge in S appears in a cycle: the corresponding triangle. In order to show that S is a cactus, we need to prove that each cycle has length three. We can substitute f, f' by e, e' and add the third edge for each pair $\{f, f'\}$ one at a time, and we maintain the invariant that each cycle consists of $\{e, e'\}$, corresponding to a matched pair $\{f, f'\}$ of F, and the third edge of the corresponding triangle. This property is true before substituting the first pair and adding any third edge, since F is a forest. Let us prove that we maintain the invariant after substituting f, f' by e, e' and adding the third edge e'' of a triangle whose edges are $\{e, e', e''\}$. We know that before substituting f, f' by e, e' and adding e'', f and f' did not appear in any cycle, by the invariant. We need to show that substituting f, f' by e, e' and adding e'' only creates the cycle e, e', e''. Because e is parallel to f and e' is parallel to f', any cycle created by the substitution of f, f' by e, e' and the addition of e'' has to contain e'' (otherwise f and f' would appear in a cycle).

Assume by contradiction that there is a cycle C which is not e, e', e''.

Assume first that C contains neither e nor e'. By replacing e'' by f and f' in C, we either get a cycle with f and f' and without e, e', e'', a contradiction, or a union of two cycles (this is the case in which C goes through the common endpoint of f and f'). But one of these cycles contains f and does not contain e, e', e'', a contradiction. If C contains e and e'' but not e', then by replacing e and e'' by f' in C we get a cycle containing f' and not e, e', e'', contradicting the invariant. The case in which C contains e' and e'' but not e is similar. It follows that S is a cactus.

For each two edges in F we get a triangle in S, for a total of p triangles. This completes the proof. ■

As described by Chiba and Nishizeki [CN85], we can explicitly list all the triangles in a graph G with m edges in time $O(m^{3/2})$. So $|E'|$ is $O(m^{3/2})$.

Gabow and Stallmann [GS85] describe an algorithm for GMP, which runs in time $O(m'n' \log^6 n')$, where m' and n' are the number of edges and vertices, respectively, in the input graph. In our case, $n' = n$ and $m' = |E'|$, which is $O(m^{3/2})$. This gives a time bound of $O(m^{3/2} n \log^6 n)$ for this phase.

From the output of the Gabow-Stallmann algorithm, it is easy to find a maximum triangular cactus in time $O(n)$ (Lemma 2.13). Therefore the total time is $O(m^{3/2} n \log^6 n)$.

3 Outerplanar Subgraphs

Serendipitously, Algorithm B produces outerplanar graphs, so it is an approximation algorithm for MAXIMUM OUTERPLANAR SUBGRAPH, which is NP-Complete [GJ79, p. 197].

In fact, any algorithm which produces a spanning tree has performance ratio at least 1/2, because any outerplanar graph on $n \geq 2$ vertices has at most $2n-3$ edges (see below). A careful analysis shows that the performance ratio of B when used for MAXIMUM OUTERPLANAR SUBGRAPH is at least 2/3. This is an easy consequence of Theorem 3.2, in order to prove which we need some preliminaries.

An outerplanar graph G is a *maximal outerplanar graph* if no edge can be added without losing outerplanarity. As mentioned in [H72, p. 106], every maximal outerplanar graph G with at least three vertices is a triangulation of a polygon (i.e., the boundary of the exterior face is a Hamiltonian cycle and each interior face is triangular). By [H72, Cor. 11.9], G must have a vertex of degree two and $2|V(G)|-3$ edges (this last statement is also true for $|V(G)| = 2$).

Lemma 3.1 *Let H be an embedding of a maximal outerplanar graph as the triangulation of a polygon. If H has an odd number $n = 2p+1$ of vertices, then there is a triangular cactus in H with p triangles. If H*



Figure 2: Graphs H_1 and H_2, built from H. The shaded region gives an example of a triangular structure of the desired size, like the one constructed in the last paragraph of the proof of Lemma 3.1.

has an even number $n = 2p$ of vertices and xy is an edge on the boundary of the exterior face, then there is a triangular cactus S in H with $p-1$ triangles such that in S, x and y are not connected.

Notice that we obtain the maximum number of triangles possible. In the former case all vertices are in the same component of the cactus, while in the latter, the cactus has two components.

Proof. We use a plane embedding of H.

The proof is by induction on n, the number of vertices of H. The case $n = 1$ is trivial. If $n = 2$ (in this case there is only one edge and $p = 1$), the theorem is true.

We inductively construct a triangular cactus of the given size.

Let $n = 2p+1$. Let u be a vertex of degree two. Let x and y be its neighbors. They are adjacent, since interior faces are triangles. The graph $H-\{u\}$ is maximal outerplanar (since it has $(2n-3)-2 = 2(n-1)-3$ edges) and has an even number of vertices. It is easy to check that if a triangular cactus S' in this smaller graph has the property that x and y are not connected in S', we can add the triangle xyu to get a triangular cactus in H. The size of this cactus is $p-1$, by induction, plus one, for a total of p.

Let $n = 2p$ and let the edge xy be on the boundary of the exterior face. This edge is on the boundary of a triangular face xyv on the inside. Walking along the Hamiltonian cycle which is the boundary of the exterior face, starting at v and in the direction that visits x just before y, let D_1 be the set of vertices visited between v and x, inclusive, and let $n_1 = |D_1|$. Walking along the Hamiltonian cycle in the opposite direction again starting at v, let D_2 be the set of vertices visited between v and y, inclusive, and let $n_2 = |D_2|$; $D_1 \cap D_2 = \{v\}$ and $D_1 \cup D_2 = V(H)$. The only edge in H between $D_1 - \{v\}$ and $D_2 - \{v\}$ is the edge xy.

Let H_1 be the subgraph of H induced by vertex set D_1, with, say, e_1 edges, and let H_2 be the subgraph of H induced by vertex set D_2, with, say, e_2 edges (see Figure 2).

We have $n_1 + n_2$ is odd, since v is counted twice. Let us say without loss of generality that $n_1 = 2p_1 + 1$ is odd and $n_2 = 2p_2$ is even. Then $n = 2(p_1 + p_2)$. We have $e_1 + e_2 = (2n-3) - 1$, as from H only the edge xy is not an edge of either H_1 or H_2. Since $e_1 \leq 2n_1 - 3$ and $e_2 \leq 2n_2 - 3$, we infer that

$e_1 + e_2 \leq 2(n_1 + n_2) - 6 = 2(n+1) - 6 = 2n - 4$. Since, in fact, $e_1 + e_2 = 2n - 4$, we infer that $e_1 = 2n_1 - 3$ and $e_2 = 2n_2 - 3$. Thus both H_1 and H_2 have to be maximal outerplanar, as they have the maximum allowable number of edges.

Then by the inductive hypothesis we can construct in H_1 a cactus S_1 with p_1 triangles. If we apply the inductive hypotheses to H_2 with vy being the edge on the exterior face, we obtain a triangular cactus S_2 with $p_2 - 1$ triangles in which y and v are not connected. Then putting together the edges of S_1 and S_2 we get S, a cactus in H. In the new cactus S, any possible $x - y$-path must visit v, since neither S_1 nor S_2 has edge xy. But in S_2, y and v are not connected. It follows that x and y are not connected in S, so S is the desired cactus. S has $p_1 + (p_2 - 1)$ triangles, which is exactly the number we wanted. ■

In conclusion, for a maximal outerplanar graph with n vertices, we can find a triangular structure with $\lfloor \frac{n-1}{2} \rfloor$ triangles.

Now we prove a lower bound on $\rho(H)$.

Theorem 3.2 *If H is outerplanar, then $\rho(H) \geq 2/3$.*

Proof. Let H be any 2-connected outerplanar graph. We add t edges to obtain a maximal outerplanar plane graph H'. Note that H' has $2n - 3$ edges and a triangular structure S with exactly $\lfloor \frac{n-1}{2} \rfloor$ triangles.

However, the t missing edges can destroy at most t of these triangles in S, because S is a cactus. If $t \geq n/2$, we infer that

$$\rho(H) \geq \frac{n-1}{2n - 3 - n/2} > \frac{2}{3}.$$

Assume to the contrary that $t \leq \lfloor \frac{n-1}{2} \rfloor$. Then the number of edges in the triangular structure S is at least $n - 1 + (\lfloor \frac{n-1}{2} \rfloor - t)$. Thus

$$\rho(H) \geq \frac{n - 1 + \lfloor \frac{n-1}{2} \rfloor - t}{2n - 3 - t}.$$

The worst case is achieved when $t = \lfloor \frac{n-1}{2} \rfloor$ and is $2/3$.

If H is not 2-connected, we can do the above analysis for each of the 2-connected components (an edge appears in exactly one 2-connected component) and infer that a maximum triangular structure has $2/3$ of the edges in H. ■

The theorem above is tight, in the sense that there are outerplanar graphs H for which $\rho(H)$ is arbitrarily close to $2/3$. In fact, there are outerplanar graphs H_i with $2i$ vertices and $3i - 2$ edges which do not have any triangle.

Corollary 3.3 *Algorithm B has performance ratio 2/3 for MAXIMUM OUTERPLANAR SUBGRAPH.*

4 The Complexity of the Problems

Papadimitriou and Yannakakis [PY91] defined a natural variant of NP for optimization problems: the complexity class Max SNP. This class, as they have shown, contains several well-known optimization problems, such as MAX 3-SAT and MAXIMUM CUT. In this section, we prove that MAXIMUM PLANAR SUBGRAPH (MPS) is Max SNP-hard, as is its complementary version: given a graph, find a smallest subset of its edges whose removal results in a planar graph. This means, by results of Arora et al. [ALMSS92], that there is a constant $\epsilon > 0$ such that the existence of a polynomial-time approximation algorithm for MPS with performance ratio at least $1 - \epsilon$ implies that $P = NP$, and that an analogous statement can be made about the complementary problem.

As in [PY91], we use the concept of *L-reduction*, which is a special kind of reduction that preserves approximability. Let A and B be two optimization problems. We say *A L-reduces to B* if there are two polynomial-time algorithms f and g, and positive constants α and β, such that for each instance I of A,



Figure 3: Graph H' constructed from H.

1. Algorithm f produces an instance $I' = f(I)$ of B, such that the optima of I and I', of costs denoted $Opt_A(I)$ and $Opt_B(I')$ respectively, satisfy $Opt_B(I') \leq \alpha \cdot Opt_A(I)$, and
2. Given any feasible solution of I' with cost c', algorithm g produces a solution of I with cost c such that $|c - Opt_A(I)| \leq \beta \cdot |c' - Opt_B(I')|$.

The main result of this section is

Theorem 4.1 *MAXIMUM PLANAR SUBGRAPH is Max SNP-hard.*

Proof. Denote by $TSP_4(1,2)$ the following variant of the traveling salesman problem: given a complete graph, a pair of distinct vertices x, y, and costs one or two for each edge, such that the graph induced by the edges of cost one has maximum degree at most four, find a Hamiltonian path from x to y of minimum cost. Papadimitriou and Yannakakis [PY93] showed that $TSP_4(1,2)$ is Max SNP-hard.

We shall prove $TSP_4(1,2)$ L-reduces to MPS. The basic idea of the reduction comes from Liu and Geldmacher [LG77], where the decision version of MPS is proved to be NP-complete.

The first part of the L-reduction is the polynomial-time algorithm f and the constant α. Given any instance I of $TSP_4(1,2)$, f produces an instance G of MPS such that the cost of the optimum of G in MPS, denoted $Opt_{MPS}(G)$, is at most α times the cost of the optimum of I in $TSP_4(1,2)$, denoted by $Opt_{TSP_4(1,2)}(I)$, i.e., $Opt_{MPS}(G) \leq \alpha \cdot Opt_{TSP_4(1,2)}(I)$.

Consider an instance I of $TSP_4(1,2)$. I is a complete graph $K = (V, E)$, a pair of distinct vertices x, y of V and a subset E_1 of E consisting of the edges of cost one. Let $H = (V, E_1)$ and $H' = (V \cup T, E_1 \cup F_1 \cup F_2)$, where $T = \{t_0, t_1, t_2, t_3\}$, $T \cap V = \emptyset$, $F_1 = \{t_0t_1, t_0t_3, t_1t_2, t_1t_3, t_1x, t_2t_3, t_3y\}$ and $F_2 = \cup_{z \in V}\{t_0z, t_2z\}$ (see Figure 3).

Denoting by n the number of vertices of H, let G be the graph obtained from H' by (1) replacing each edge e in F_1 by $2n$ parallel internally-disjoint paths of length two (having new internal vertices) between the endpoints of e and (2) replacing each edge e in F_2 by eight parallel internally-disjoint paths of length two (having new internal vertices) between the endpoints of e (see Figure 4).

Clearly G can be obtained from I in time polynomial in the size of I.



Figure 4: Edge e and its substitution by internally-disjoint paths of length two.

Lemma 4.2 $Opt_{MPS}(G) \leq 124 \cdot Opt_{TSP_4(1,2)}(I)$.

Proof. Observe that $Opt_{TSP_4(1,2)}(I) \geq n-1$. A clear upper bound for $Opt_{MPS}(G)$ is the number of edges of G. To compute this, note first that H has maximum degree at most four by the definition of $TSP_4(1,2)$, and so H has at most $2n$ edges. Let us call the edges in F_1 *2n-edges* and the edges in F_2 *8-edges*. There are seven $2n$-edges: F_1 contains seven edges, each of them corresponding to $4n$ edges in G. There are $2n$ 8-edges: F_2 contains $2n$ edges, each of them corresponding to 16 edges in G. Hence the number of edges in G outside of H is $7 \cdot 4n + 16 \cdot 2n = 60n$. The total number of edges in G is therefore at most $2n + 60n = 62n \leq 124(n-1)$. Therefore $Opt_{MPS}(G) \leq 124(n-1) \leq 124 \cdot Opt_{TSP_4(1,2)}(I)$. ■

This finishes the first part of the L-reduction, since we can take $\alpha = 124$.

The second and hard part of the L-reduction is the constant β and the algorithm g. Given a planar subgraph of G with m edges, g produces in polynomial time a Hamiltonian path from x to y of cost t in K such that $|t - Opt_{TSP_4(1,2)}(I)| \leq \beta |m - Opt_{MPS}(G)|$. We shall see that $\beta = 1$ suffices.

First, given a planar subgraph P of G, let us describe another planar subgraph P' of G with at least as many edges as P. Moreover, P' shall contain all edges of G not in H.

Let e be a $2n$-edge or an 8-edge of H'. We say e *appears in* P if P contains both edges of *all* the paths of length two corresponding to e. In this case, we also say that the endpoints of e *are adjacent in* P *by the* $2n$*-edge* or *8-edge* e. We say e *is missing in* P if P contains both edges of *none* of the paths (of length two) corresponding to e. (In this case, if e is an 8-edge, then P is missing at least eight of its 16 edges in G corresponding to e.) It is possible that a $2n$-edge or an 8-edge of H' neither appears in P nor is missing in P.

Let us modify P so that any $2n$-edge or 8-edge of H' either appears in P or is missing in P. This is done as follows. If a $2n$-edge or an 8-edge e of H' neither appears nor is missing in P, then we insert in P all edges of G in the paths (of length two) corresponding to e. Note that P remains planar. Clearly, the number of edges in P cannot decrease by this operation. The new graph is also called P.

Now we can describe P'. We have three cases: (1) if some $2n$-edge e does not appear in P, then define P' to be the graph induced by all edges of G not in H; (2) if all the $2n$-edges and the 8-edges of H' appear in P, then let P' be the same as P; and (3) if all the $2n$-edges of H' appear in P, but not all the 8-edges, then we modify P to obtain P', as described in the next two paragraphs.

The idea is to remove from P some edges of H and add to P edges of H' not in H so that all the 8-edges of H' appear in the modified graph, and it remains planar and has at least as many edges as the original P.

Let U be the set of vertices v of H such that at least one of the two 8-edges incident to v in H' is missing in P. Observe that $|U| \geq 1$, as case (2) considered $|U| = 0$. For each vertex v in U, remove from P all edges of H incident to v in P (at most four edges are removed per vertex) and add to P all



Figure 5: Cycle C, regions R_1 and $R_2 = R_2' \cup R_2''$.

the edges outside of H so that the two 8-edges incident to v appear in P (at least eight edges are added, corresponding to the 8-edges incident to v missing in P). To guarantee that the graph obtained this way is planar, we must make room to embed the modified 8-edges. This is done by also removing from P all edges of H incident to y (if they were not already removed). Let P' be the graph obtained after all these modifications.

Lemma 4.3 *P' is planar and has at least as many edges as P.*

Proof. In case (1), we include in P' at least $2n$ edges that do not appear in P (at least one in each of the $2n$ paths corresponding to e), and we remove at most $2n$ edges, the maximum number of edges in H. So P' has at least as many edges as P. Moreover P' is planar.

There is nothing to be proved in case (2).

Case (3) is the complicated one. First note that P' has at least as many edges as G, since, for each vertex in U, we remove at most four edges and add at least eight. Furthermore, we remove at most four edges incident to y. Hence we gain at least $(8-4)|U| - 4 = 4|U| - 4 \geq 0$ edges, since $|U| \geq 1$.

Now, let us show that P' is planar. We can think of the $2n$-edges and 8-edges as single edges, as they are in H' (since if we can embed a single edge, we can embed a $2n$-edge or an 8-edge as well). We will modify a given embedding of P into an embedding for P'.

Let C be the cycle (using four $2n$-edges) t_0, t_1, t_2, t_3, t_0. Observe that the $2n$-edges in C appear in P, since we are in case (3). Given an embedding for P, cycle C divides the plane into two regions, R_1, containing the $2n$-edge t_1t_3, and R_2 (see Figure 5). The $2n$-edge t_1t_3 separates t_0 from t_2 in R_1. Moreover, each vertex in $V - U$ (the vertices of H not in U) is adjacent in P by 8-edges to t_0 and t_2. Because t_0 and t_2 are separated in R_1, none of these vertices can be embedded in R_1, which implies they must be embedded in R_2. Keep these vertices (t_0, t_1, t_2, t_3 and the vertices in $V - U$) embedded as they are.

Now, observe that y is adjacent in P' only to t_0, t_2 (by 8-edges) and t_3 (by a $2n$-edge). Furthermore, y is the only vertex in H which is adjacent to t_3. This means, before we embed y, the vertices t_0, t_2 and t_3 are not separated in R_2. Therefore y can be embedded in R_2 with the $2n$-edge t_3y and the 8-edges t_0y and yt_2 "next to" the $2n$-edges t_0t_3 and t_3t_2. The edges t_0y and yt_2 together split region R_2 into two regions R_2' containing edge t_3y, and R_2'' containing vertex t_1. Observe that t_0 and t_2 are not separated in R_2'', since t_3 is the only vertex, besides t_0 and t_2, which is adjacent to y (by a $2n$-edge).

All the vertices in $U - \{x\}$ are adjacent in P' only to t_0 and t_2 (by 8-edges), and therefore they all can be embedded in R_2'' with their two 8-edges "next to" the 8-edges t_0y and yt_2. If $x \in U$ then observe that x is the only vertex in H which is adjacent in P' to t_1 (by a $2n$-edge). This means, before we embed x,

the vertices t_0, t_1 and t_2 are not separated in R_2''. Therefore x can be embedded in R_2'' with the $2n$-edge xt_1, and the 8-edges t_0x and xt_2 "next to" the $2n$-edges t_0t_1 and t_1t_2. If $x \notin U$ then it does not need to be moved in the embedding. The embedding obtained this way is a plane embedding of P', completing the proof that P' is planar. ■

Observe that P' contains all the edges of G not in H. Let F be the set of edges of H appearing in P'.

Lemma 4.4 *The graph $G_F = (V, F)$ is a collection of vertex-disjoint paths which can be extended in K (the complete graph on V) to a Hamiltonian path from x to y, in polynomial time.*

Proof. Let us prove that G_F satisfies the following four conditions: (1) There is no vertex of degree greater than two in G_F. (2) Vertices x and y have degree at most one in G_F. (3) There is no cycle in G_F. (4) If x and y are in the same component of G_F, then this component spans all vertices in V. We will prove each of these conditions holds by contradiction.

Suppose (1) does not hold. Let z_0 be a vertex in V of degree at least 3 in G_F. Let z_1, z_2, z_3 be three of its neighbors in G_F. (Notice that z_0, z_1, z_2, z_3 are distinct vertices of H, so they are distinct of t_0, t_2.) Then each one of t_0, t_2, z_0 is adjacent in P' to each one of z_1, z_2, z_3 (some of them are adjacent in P' by 8-edges). Therefore $t_0, t_2, z_0; z_1, z_2, z_3$ define a subdivision of $K_{3,3}$ in P', a contradiction, because P' is planar. Thus (1) holds.

Suppose (2) does not hold. If x has degree more than one in G_F, let z_1 and z_2 be two of its neighbors in G_F. (Notice that z_1 and z_2 are distinct vertices of H, distinct of x, so they are distinct of t_0, t_2.) Then each one of t_0, t_2, x is adjacent in P' to each one of t_1, z_1, z_2 (some of them are adjacent in P' by $2n$-edges or 8-edges). Therefore $t_0, t_2, x; t_1, z_1, z_2$ define a subdivision of $K_{3,3}$ in P', a contradiction, because P' is planar. Analogously, we have a contradiction if y has degree more than one in G_F. Thus (2) holds.

Suppose (3) does not hold. Let z_1, z_2, z_3 be three vertices in a cycle of G_F. (Since z_1, z_2, z_3 must have degree at least two, they are not x or y by condition (2), and they are not t_0 or t_2 since they are vertices of H.) Then z_1, z_2, z_3, t_0, t_2 are pairwise linked by internally vertex-disjoint paths (the path between t_0 and t_2 uses the two 8-edges incident to x, while the others use one $2n$-edge or 8-edge). Therefore t_0, t_2, z_1, z_2, z_3 define a subdivision of K_5 in P', a contradiction, because P' is planar. Hence (3) holds.

Suppose (4) does not hold. Let z_0 be a vertex in V which is not in the component having x and y in G_F. In this case, t_0, t_1, t_2, x, y are pairwise linked by internally vertex-disjoint paths (the path between t_0 and t_2 uses z_0, the path between t_1 and y uses t_3, the path between x and y is in G_F, and the others use one $2n$-edge or 8-edge). Therefore t_0, t_1, t_2, x, y define a subdivision of K_5 in P', a contradiction, because P' is planar. Hence (4) holds.

Therefore the conditions hold. From (1) and (2), we conclude that G_F is a collection of paths. From (2) and (3), these paths can be extended in K (the complete graph on V) to a Hamiltonian path from x to y. Furthermore, note that this can be done in polynomial time. ■

Let HP be a Hamiltonian path from x to y containing all edges in F and some edges (in K) of cost two. HP exists by Lemma 4.4. Denote by m' the number of edges of P' and by t the cost of HP.

Now the following Lemma states that β exists, and specifically, $\beta = 1$. This will complete the proof of the theorem.

Lemma 4.5

$$t - Opt_{TSP_4(1,2)}(I) = Opt_{MPS}(G) - m',$$

and hence

$$|t - Opt_{TSP_4(1,2)}(I)| = |m' - Opt_{MPS}(G)|.$$

Proof. As in the proof of Lemma 4.2, the number of edges in G outside of H is $60n$. All these edges are in P'. Therefore the number of edges in F is $m' - 60n$. And the cost of HP is

$$t = 2(n-1) - (m' - 60n). \tag{9}$$

Let Q be an optimal solution of MPS for G. Using the same argument for Q that we used for P, Lemma 4.4 and the argument above imply the existence of a Hamiltonian path of cost $2(n-1)-(Opt_{MPS}(G)-60n)$. Therefore

$$Opt_{TSP_4(1,2)}(I) \leq 2(n-1) - (Opt_{MPS}(G) - 60n). \tag{10}$$

Given an optimum solution HP^* of $TSP_4(1,2)$ for I, we can construct a solution of MPS for G by selecting all the edges outside of H plus the edges of cost one in HP^*. Observe that these edges really determine a planar subgraph of G. Let z be the number of these edges. Since we have $60n$ edges in G outside of H, HP^* has $z-60n$ edges of cost one and the remaining ones (of its $n-1$ edges) have cost two. This means that

$$\begin{aligned} Opt_{TSP_4(1,2)}(I) &= 2(n-1) - (z - 60n) \\ &\geq 2(n-1) - (Opt_{MPS}(G) - 60n), \end{aligned}$$

since $Opt_{MPS}(G) \geq z$.

Therefore, from (10) and (11), we have

$$Opt_{TSP_4(1,2)}(I) = 2(n-1) - (Opt_{MPS}(G) - 60n).$$

And this together with (9) means

$$t - Opt_{TSP_4(1,2)}(I) = Opt_{MPS}(G) - m'.$$

Hence $|t - Opt_{TSP_4(1,2)}(I)| = |m' - Opt_{MPS}(G)|$. ■

From $m \leq m'$, it follows that

$$t - Opt_{TSP_4(1,2)}(I) \leq Opt_{MPS}(G) - m$$

and

$$|t - Opt_{TSP_4(1,2)}(I)| \leq 1 \cdot |m - Opt_{MPS}(G)|.$$

This completes the proof of Theorem 4.1. ■

Let us denote the complementary version of MPS by NPD: given a graph G, find a smallest set of edges of G whose removal results in a planar graph.

A slight modification of the L-reduction presented above proves the following.

Theorem 4.6 *NPD is Max SNP-hard.*

Proof. The first part of the L-reduction is almost the same. From an instance I of $TSP_4(1,2)$, we construct G in exactly the same way. As before, $Opt_{TSP_4(1,2)}(I) \geq n-1$. As in the proof of Lemma 4.2, the maximum number of edges of G is $62n$. Thus the optimum of G in NPD, denoted as $Opt_{NPD}(G)$, is at most $62n$. And then $Opt_{NPD}(G) \leq 62n \leq 124(n-1) \leq 124 \cdot Opt_{TSP_4(1,2)}(I)$. We can take $\alpha = 124$, as before.

In the second part, given an instance I of $TSP_4(1,2)$, let G be constructed from I as in the previous reduction. Let D be a subset of the edges of G whose removal results in a planar subgraph of G. We shall find in polynomial time a Hamiltonian path HP such that $t - Opt_{TSP_4(1,2)}(I) \leq d - Opt_{NPD}(G)$, where t

is the number of edges in HP and $d = |D|$. Just as we took a planar subgraph P of G and found a planar subgraph P' which contains all edges of G not in H, and is at least as large as P, from D we can find a set D' of edges of G containing *none* of the edges of G not in H, which is at least as *small* as D. Applying Lemma 4.4, we can obtain a Hamiltonian path HP, as before, in polynomial time.

Now, let us prove that β exists. Let m' be the number of edges in P'. Note that $d' + m' = |E(G)|$. Moreover, $Opt_{MPS}(G) + Opt_{NPD}(G) = |E(G)|$. Therefore $d' - Opt_{NPD}(G) = Opt_{MPS}(G) - m'$. Applying Lemma 4.5, we conclude that $t - Opt_{TSP_4(1,2)}(I) = d' - Opt_{NPD}(G)$, which, together with $d' \leq d$, implies that we can take $\beta = 1$. ■

5 Open Problems

Many open problems are suggested by this research. How large a performance ratio can one achieve is an obvious one. Is there a linear-time approximation algorithm for MAXIMUM PLANAR SUBGRAPH with performance ratio $1/3 + \epsilon$? (A *maximal* planar subgraph can be found in linear time [H95, D95].) Is there any approximation algorithm with a constant performance ratio for NPD? What performance ratio can be achieved for THICKNESS (given G, partition the edges of G into as few planar subgraphs as possible)? A factor of 3 here is trivial, via arboricity.

6 Acknowledgments

The authors are very grateful to David Williamson for his help. We thank Laci Lovász for informing us that a maximum triangular structure can be found in polynomial time. We thank Robert Cimikowski for kindly sending us drafts of his papers. The third author would like to thank Kurt Mehlhorn, for suggesting the problem and supporting his work. We are also very grateful to Robin Thomas for helpful discussions, and specifically for providing a simpler proof of Claim 2.8. We also would like to thank the program committee of SODA'96 for informing us about a faster way to find all triangles in a graph.

References

[ALMSS92] S. Arora, C. Lund, R. Motwani, M. Sudan and M. Szegedy, "Proof Verification and Hardness of Approximation Problems," *Proc.* 33^{rd} *IEEE Symposium on Foundations of Computer Science*, 14–23, 1992.

[BM76] J. A. Bondy and U. S. R. Murty, *Graph Theory with Applications*, MacMillan Press, 1976.

[CHT93] J. Cai, X. Han and R. E. Tarjan, "An $O(m \log n)$-time Algorithm for the Maximal Planar Subgraph Problem," *SIAM Journal on Computing*, 22:1142–1162, 1993.

[Cim95] R. Cimikowski. "An Analysis of Some Heuristics for the Maximum Planar Subgraph Problem," *Proc.* 6^{th} *Annual ACM-SIAM Symp. on Discrete Algorithms*, 322–331, 1995.

[CNS79] T. Chiba, I. Nishioka and I. Shirakawa. "An Algorithm of Maximal Planarization of Graphs," *Proc. IEEE Symp. on Circuits and Systems*, 649–652, 1979.

[CN85] N. Chiba and T. Nishizeki, "Arboricity and Subgraph Listing Algorithms," *SIAM Journal of Computing*, 14:210-223, 1985.

[D95] H. N. Djidjev, "A Linear Algorithm for the Maximal Planar Subgraph Problem," *Proc.* 4^{th} *International Workshop on Algorithms and Data Structures* (WADS '95), 369-380, 1995.

[DFF85] M. E. Dyer, L. R. Foulds and A. M. Frieze, "Analysis of Heuristics for Finding a Maximum Weight Planar Subgraph," *European Journal of Operational Research*, 20:102–114, 1985.

[F92] L. R. Foulds, *Graph Theory Applications*, Springer-Verlag, New York, 1992.

[GS85] H. N. Gabow and M. Stallmann, "Efficient Algorithms for Graphic Matroid Intersection and Parity," *Automata, Language and Programming:* 12^{th} *Colloq., Lecture Notes in Computer Science*, Vol. 194, 210–220, 1985.

[GJ79] M. R. Garey and D. S. Johnson, *Computers and Intractability*, W. H. Freeman and Co., 1979.

[H72] F. Harary, *Graph Theory*, Addison-Wesley Publishing Company, 1972.

[H95] W. Hsu, "A Linear Time Algorithm for Finding Maximal Planar Subgraphs," *Proc.* 6^{th} *Annual International Symposium on Algorithms and Computation* (ISAAC95), 1995.

[JM93] M. Jünger and P. Mutzel, "Maximum Planar Subgraph and Nice Embeddings: Practical Layout Tools," *Universität zu Köln, Report No. 93.145*, 1993, to appear in *Special Issue of Algorithmica on Graph Drawing.*

[LG77] P. C. Liu and R. C. Geldmacher, "On the Deletion of Nonplanar Edges of a Graph," *Proc.* 10^{th} *Southeastern Conference on Combinatorics, Graph Theory, and Computing*, 727–738, 1977.

[LP86] L. Lovász and M. D. Plummer, *Matching Theory*, Elsevier Science, Amsterdam, 1986.

[PY91] C. H. Papadimitriou and M. Yannakakis. "Optimization, Approximation, and Complexity Classes," *Journal of Computer and System Sciences*, 43:425–440, 1991.

[PY93] C. H. Papadimitriou and M. Yannakakis, "The Traveling Salesman Problem with Distances One and Two," *Mathematics of Operations Research*, 18(1):1–11, 1993.

[TDB88] R. Tamassia, G. Di Battista and C. Batini, "Automatic Graph Drawing and Readability of Diagrams," *IEEE Transactions on Systems, Man and Cybernetics*, 18:61-79, 1988.